

A Nonlinear Approximation of Operator Equation $V^*QV = Q$: Nonspectral Decomposition of Nonnormal Operator and Theory of Stability

S.A. Chorošavin

Abstract

V denotes arbitrary bounded bijection on Hilbert space H . We try to describe the sets of V -stable vectors, i.e. $\{x \in H \mid \text{the sequence } \|V^N x\| (N = 1, 2, \dots) \text{ is bounded}\}$ and some other analogous sets. We do it in terms of one-parameter operator equation $Q = V^*(Q + tI)(I + tQ)^{-1}V$ (t is real valued parameter $0 \leq t \leq 1$, Q is operator to be found, $0 \leq Q$)

1 Introduction

Throughout this paper H will denote a Hilbert space with scalar product $\langle, \rangle$, V denotes a linear bounded bijection H onto H ,

$$r(T) := \text{spectral radius of } T.$$

We will discuss the structure of the next four sets:

$$Stab_+(V) := \{x \in H \mid \forall a > 1 \exists C \geq 0 \forall N \geq 0 \quad \|V^N x\| \leq Ca^N\}.$$

$$Stab(V) := \{x \in H \mid \|V^n x\| \leq M \text{ for some real } M \text{ and every } n = 0, 1, 2, \dots\}$$

$$Stab_0(V) := \{x \in H \mid \|V^n x\| \rightarrow 0 \quad (n \rightarrow \infty)\}$$

$$l_2(V) := \{x \in H \mid \|Vx\|^2 + \|V^2x\|^2 + \dots + \|V^n x\|^2 + \dots < \infty\}$$

Recall V is **similar to an unitary operator** iff there exists a bounded uniformly positive operator Q such that

$$V^*QV = Q$$

With this equation we shall consider an 'approximation' equation (parametrized by real t)

$$Q = V^* \frac{Q + t}{I + tQ} V, \quad Q \geq 0, \quad 0 < t \leq 1 \quad (*)$$

(hereinafter t denotes always a real number such that $0 < t \leq 1$ and if no confusion can occur we shall often write t instead of tI , I is identity operator).

The interest in this equation can be motivated by the next

Example 1. Let V be normal. Routine, though tedious calculation shows that

$$Q_t := \left[-\frac{I - V^*V}{2} + \sqrt{\left(\frac{I - V^*V}{2}\right)^2 + t^2 V^*V} \right] \frac{1}{t}$$

is uniformly positive solution of (*), there holds

$$Q_t^{-1} = \left[-\frac{I - (V^*V)^{-1}}{2} + \sqrt{\left(\frac{I - (V^*V)^{-1}}{2} \right)^2 + t^2(V^*V)^{-1}} \right] \frac{1}{t}$$

and there exist

$$X_0 := \text{strong} - \lim tQ_t = (V^*V - I)E(1, \infty) \quad (t \rightarrow +0)$$

$$Y_0 := \text{strong} - \lim tQ_t^{-1} = ((V^*V)^{-1} - 1)E(0, 1) \quad (t \rightarrow +0)$$

Besides that it is fast evident that $Q_tE([1]) = E([1])$, $Q_tE(1, \infty)$ is monotone increasing (with $t \rightarrow +0$), $Q_tE(0, 1)$ is monotone decreasing and there exists an

$$R_0 := \text{strong} - \lim (I + Q_t)^{-1} = E(0, 1) + E([1])/2 \quad (t \rightarrow +0).$$

here $E(\Delta)$ denotes the spectral function of the selfadjoint operator V^*V . Note:

$$I - R_0 = E([1])/2 + E(1, \infty)$$

$$\text{Ker}(I - R_0) = E(0, 1)H$$

$$\text{Ran}X_0 \subset E(1, \infty)H = \overline{\text{Ran}X_0}$$

$$\text{Ran}Y_0 \subset E(0, 1)H = \overline{\text{Ran}Y_0}$$

□

Also, in the above considered case of the normal V it is established that the operators X_0, Y_0, R_0 define (in essential) the spectral subspaces of V (with V together one can consider $aV - b, b/a \notin \text{spectrum } V$). In this article we shall show that the similar situation holds for the arbitrary bounded bijection V .

We follow standards of [RS] when we apply mathematical concepts and sometimes we apply P.A.M. Dirac's 'bra-ket' syntax.

We will often cite some assertions and propositions of [Ch1-4]. For the most convenient and accesible way to do it, we collect them together and resume them here as

Theorem 1.

- (i) *the solution of (*) exists and it is unique; denote it by Q_t*
(ii) *Q_t is bounded selfadjoint uniformly positive and there are satisfied inequalities:*

$$tV^*V \leq Q_t \leq V^*V/t$$

$$tQ_t \leq sQ_s \quad (0 < t \leq s \leq 1)$$

denote $X_t := tQ_t$

- (iii) *Q_t^{-1} is (unique) solution of the analogous equation:*

$$Q_t^{-1} = V^{-1} \frac{Q_t^{-1} + t}{I + tQ_t^{-1}} (V^{-1})^*$$

so there are satisfied inequalities:

$$tV^{-1}(V^{-1})^* \leq Q_t^{-1} \leq V^{-1}(V^{-1})^*/t$$

$$tQ_t^{-1} \leq sQ_s^{-1} \quad (0 < t \leq s \leq 1)$$

denote $Y_t := tQ_t^{-1}$.

- (iv) *Let*

$$X_0 := \text{strong} - \lim X_t, \quad Y_0 := \text{strong} - \lim Y_t \quad (t \rightarrow +0)$$

The operators X_0, Y_0 are bounded positive and they are maximal solutions of the equations

$$X = V^* \frac{X}{I + X} V, \quad X \geq 0, \quad \text{resp.} \quad VYV^* = \frac{Y}{I + Y}, \quad Y \geq 0$$

('maximal' denotes 'maximal with respect to usual partial order for bounded operators on Hilbert space')

- (v) *There hold the formulae*

$$Y_0 = \text{strong} - \lim (V^*V + V^{*2}V^2 + \dots + V^{*n}V^n)^{-1}, \quad (n \rightarrow \infty)$$

$$X_0 = \text{strong} - \lim ((V^*V)^{-1} + (V^{*2}V^2)^{-1} + \dots + (V^{*n}V^n)^{-1})^{-1}, \quad (n \rightarrow \infty)$$

- (vi) *Denote $R_t := (I + Q_t)^{-1}$. Then $0 \leq R_t \leq I$, $Q_t = R_t^{-1} - I$ and the equation (*) is equivalent to the equation*

$$[R_t + t(I - R_t)](V^{-1})^*(I - R_t) = [(I - R_t) + tR_t]VR_t$$

- (vii) *Let R_0 be a weak operator limit point of the net $\{R_t, t \rightarrow +0\}$ (it is clear that R_0 exists and $0 \leq R_0 \leq I$). Then*

$$V \text{Ker} (I - R_0) = \text{Ker} (I - R_0)$$

$$V^{*-1} \text{Ker} R_0 = \text{Ker} R_0$$

In particular,

$$V \overline{\text{Ran} R_0} = \overline{\text{Ran} R_0} \quad \square$$

2 Equation $Q = V^* \frac{Q + t}{I + tQ} V$. General Properties.

Hereinafter $\mathbf{F}$ denotes an arbitrary ultra filter, which majorizes usual convergence to $+0$. We will write

$$t \rightarrow +0 \quad \text{instead of } t \xrightarrow{\mathbf{F}} +0$$

if no confusion can occur.

Definition 1.

$Fin Q := \{x \in \mathbf{H} \mid \langle x, Q_t x \rangle \leq M_x \text{ for some real } M_x \text{ and for almost every } t \text{ resp. } \mathbf{F}\}$

$$Ker Q_0 := \{x \in \mathbf{H} \mid \langle x, Q_t x \rangle \rightarrow 0 \text{ for } t \rightarrow +0\}$$

Theorem 1.

$$Ran Y_0^{1/2} = l_2(V) \subset Stab_0(V) \subset Ker Q_0 \subset Ker (I - R_0) \subset \overline{Ran R_0} \subset Ker X_0 \quad (*)$$

$$Stab(V) \subset Fin Q \subset \overline{Ran R_0} \subset Ker X_0 \quad (**)$$

Every set of these series is V -surinvariant. (recall, some L is said to be T -surinvariant, iff $TL = L$).

In addition

$$Fin Q \subset Ran R_0^{1/2}$$

Observation 0. It is well-known and evident that

$$A^*A \leq B^*B \iff \|Ax\| \leq \|Bx\| \quad (x \in H) \implies Ran A^* \subset Ran B^* \\ (A \text{ and } B \text{ are bounded})$$

Corollary 0. Let $x \in H$, Y be selfadjoint and let $Y \geq 0$. Then

$$x \in Ran Y^{1/2} \iff x\langle x \rangle \leq cY \quad \text{for some real } c.$$

Proof.

Proof of $x\langle x \rangle \leq cY \implies x \in Ran Y^{1/2}$:

Take into account Observation 0. Then obtain

$$x\langle x \rangle \leq cY \implies Ran x\langle x \rangle \subset Ran Y^{1/2} \implies x \in Ran Y^{1/2}.$$

Now proof of $x \in Ran Y^{1/2} \implies x\langle x \rangle \leq cY$:

By the definition of Ran

$$x \in Ran Y^{1/2} \iff x = Y^{1/2}y \quad \text{for an } y.$$

Hence

$$x \in Ran Y^{1/2} \implies x\langle x \rangle \equiv Y^{1/2}y\langle Y^{1/2}y \rangle = Y^{1/2}y\langle y \rangle Y^{1/2} \leq \|y\|^2 Y.$$

Now denote $\|y\|^2$ by c .

□

Observation 1. Let $\{A_t\}_t$ be a net of selfadjoint positive bounded operators. Suppose $A_t \leq aI$ for some positive number a (and every t), A_0 be a weak operator limit point of this net (clear: A_0 exists and $0 \leq A_0 \leq aI$).

Then

$$\langle x, A_t x \rangle \rightarrow 0 \Leftrightarrow \|A_t x\| \rightarrow 0 \Leftrightarrow x \in \text{Ker } A_0$$

Proof. Clear (see e.g. [Ch1]). □

Observation 2. Let Q be bounded selfadjoint positive, $0 < t \leq 1$. Then

$$\frac{Q}{I+t\|Q\|} \leq \frac{Q+t}{I+tQ} \leq Q+t$$

Corollary.

$$\boxed{V \text{Fin } Q = \text{Fin } Q, \quad V \text{Ker } Q_0 = \text{Ker } Q_0.}$$

Observation 2'. Let Q be bounded selfadjoint positive, $0 < t \leq 1$. Then

$$\frac{1-t}{1+t}Q + \frac{2t}{1+t} - \frac{Q+t}{I+tQ} = \frac{t(1-t)}{1+t} \frac{(Q-I)^2}{I+tQ}$$

In particular

$$\frac{Q+t}{I+tQ} \leq \frac{1-t}{1+t}Q + \frac{2t}{1+t}$$

Denote $z := (1-t)/(1+t)$. Then $1-z = 2t/(1+t)$ and with these denotations

$$\frac{Q+t}{I+tQ} \leq zQ + (1-z)$$

Note

$$t \rightarrow +0 \iff z \rightarrow 1-0.$$

Proof. Clear. □

Observation 3 . For Q_t the just mentioned inequality gives

$$Q_t \leq V^*(zQ_t + (1-z))V$$

and with iterating this inequality one can obtain

$$Q_t \leq (1-z)[V^*V + zV^{*2}V^2 + \dots + z^{n-1}V^{*n}V^n] + z^n V^{*n}Q_t V^n$$

In particular, given numbers z, M_0 and an $x \in H$ such that

$$0 < z < 1, \quad M_0 \geq 0, \quad \|V^n x\|^2 \leq M_0 \quad (n = 1, 2, 3, \dots)$$

then

$$\langle x, Q_t x \rangle \leq (1-z)(\|Vx\|^2 + z\|V^2x\|^2 + z^2\|V^3x\|^2 + \dots) \leq M_0.$$

In particular

$$\boxed{\text{Stab}(V) \subset \text{Fin } Q}$$

Now suppose $\|V^n x\| \rightarrow 0$ ($n \rightarrow \infty$) for some $x \in H$. Let $t \rightarrow +0$. Then $z \rightarrow 1-0$ and hence $\langle x, Q_t x \rangle \rightarrow 0$ since $\|V^n x\| \rightarrow 0$ and $\langle x, Q_t x \rangle \geq 0$.

In particular

$$\boxed{\text{Stab}_0(V) \subset \text{Ker } Q_0}$$

□

Observation 4. *Let $x \in H$, $\|x\| = 1$. Then*

a)

$$x \in \text{Fin } Q \Rightarrow \exists M \geq 0 \forall \alpha \geq 0 : \quad x\langle x | \leq (M + \alpha)(Q_t + \alpha)^{-1}$$

b) Given some $M, \alpha \geq 0$, such that

$$x\langle x | \leq (M + \alpha)(Q_t + \alpha)^{-1}$$

then

$$x \in \text{Fin } Q$$

Proof

$$\begin{aligned} < x, Q_t x > \leq M & \iff < x, (Q_t + \alpha)x > \leq M + \alpha \\ & \iff \|(Q_t + \alpha)^{1/2}x\|^2 \leq M + \alpha \\ & \iff (Q_t + \alpha)^{1/2}x\langle (Q_t + \alpha)^{1/2}x | \leq M + \alpha \\ & \iff (Q_t + \alpha)^{1/2}x\langle x | (Q_t + \alpha)^{1/2} \leq M + \alpha \\ & \iff x\langle x | \leq (M + \alpha)(Q_t + \alpha)^{-1} \end{aligned}$$

□

Corollary. $\boxed{\text{Fin } Q \subset \text{Ran } R_0^{1/2} \subset \overline{\text{Ran } R_0}}$

Proof First apply Observation 4 for $\alpha := 1$ and let $t \rightarrow +0$. Then obtain

$$x\langle x | \leq (M + 1)R_0;$$

Now apply Observation 0 or Corollary 0.

□

Observation 5. *Let $x \in H$. Then*

$$x \in \text{Ran } Y_0^{1/2} \iff x\langle x | \leq cY_0 \iff < x, Q_t x > \leq ct$$

Proof

$$x \in \text{Ran } Y_0^{1/2} \iff x = Y_0^{1/2}y \implies x\langle x | \equiv Y_0^{1/2}y\langle Y_0^{1/2}y | \leq \|y\|^2 Y_0$$

Besides,

$$x\langle x | \leq cY_0 \implies \text{Ran } x\langle x | \subset \text{Ran } Y_0^{1/2} \implies x \in \text{Ran } Y_0^{1/2}$$

Now recall that

$$Y_0 \leq Y_t \text{ und } Y_t \xrightarrow{s} Y_0.$$

Hence

$$\begin{aligned} x \in \text{Ran } Y_0^{1/2} & \iff x\langle x | \leq cY_0 \\ & \iff x\langle x | \leq cY_t \\ & \iff x\langle x | \leq ctQ_t^{-1} \\ & \iff Q_t^{1/2}x\langle Q_t^{1/2}x | \leq ct \\ & \iff \|Q_t^{1/2}x\|^{1/2} \leq ct \\ & \iff < x, Q_t x > \leq ct \end{aligned}$$

□

Proof of $\boxed{Ker Q_0 \subset Ker(I - R_0)}.$

Note

$$I - R_t = Q_t(I + Q_t)^{-1} \leq Q_t$$

Now suppose $x \in Ker Q_0$ i.e. $\langle x, Q_t x \rangle \rightarrow 0$ and apply the Observation 1 to $A_t = I - R_t$.

□

Proof of $\boxed{Ker(I - R_0) \subset \overline{Ran R_0}}.$

Note Ker is closed and use definitions of Ker , Ran .

□

Proof of $\boxed{\overline{Ran R_0} \subset Ker X_0}.$

Note

$$R_t X_t = t(I + Q_t)^{-1} = X_t R_t$$

Then

$$R_t X_t = X_t R_t \xrightarrow{\|\cdot\|} 0.$$

Recall R_t, X_t are selfadjoint, positive, bounded and $X_t \xrightarrow{s} X_0$, $R_t \xrightarrow{w} R_0$. Hence $R_0 X_0 = 0 = X_0 R_0$.

The rest is evident.

□

Proof of $\boxed{Ran Y_0^{1/2} \subset l_2(V)}.$

We have

$$VY_0V^* = \frac{Y_0}{I + Y_0} \quad (*)$$

Hence $\|Y_0^{1/2}V^*x\| = \|(I + Y_0)^{1/2}Y_0^{1/2}x\| \leq \|Y_0^{1/2}x\|$ and the relation

$$W : Y_0^{1/2}x \rightarrow Y_0^{1/2}V^*x$$

defines a contraction $\overline{Ran Y_0^{1/2}} \rightarrow \overline{Ran Y_0^{1/2}}$. This contraction has extension to a contraction $\overline{Ran Y_0^{1/2}} \rightarrow \overline{Ran Y_0^{1/2}}$. It will be denote by W too. Clear, there hold

$$(i) WY_0^{1/2} = Y_0^{1/2}V^* \quad (iii) W^*W = (I + Y_0)^{-1}\overline{Ran Y_0^{1/2}}$$

$$(ii) Y_0^{1/2}W^*P = VY_0^{1/2} \quad (iv) Y_0P = ((W^*W)^{-1} - I)P$$

and $WP = PWP$, $W^*P = PW^*P$. Here is

$$P := \text{orthoprojection onto } \overline{Ran Y_0^{1/2}}$$

Hence

$$\begin{aligned} Y_0^{1/2}V^{*n}V^nY_0^{1/2} &= W^nY_0W^{*n}P = PW^nY_0W^{*n}P \\ &= PW^n((W^*W)^{-1} - I)W^{*n}P \\ &= P(W^{n-1}W^{*n-1} - W^nW^{*n})P. \end{aligned}$$

and

$$\sum_{n=2}^N \|V^nY_0^{1/2}x\|^2 = \|WPx\|^2 - \|W^N Px\|^2$$

Hence

$$\sum_0^\infty \|V^nY_0^{1/2}x\|^2 < \infty \quad \square$$

$\square$

Remark to this proof.

It follows from (*) and $Y_0 \geq 0$ that

$$VRan Y_0 = Ran Y_0, \quad V^*Ker Y_0 = Ker Y_0, \quad V\overline{Ran Y_0} = \overline{Ran Y_0}$$

Besides that,

$$(iii) \Rightarrow \|Wx\| \geq \|x\|/(1 + \|Y_0\|) \quad (x \in D_W)$$

$$(ii) \Rightarrow VRan Y_0^{1/2} \subset Ran Y_0^{1/2}$$

$$(i, \text{ or definition of } W) \Rightarrow WRan Y_0^{1/2} = Ran Y_0^{1/2} \quad \square$$

Proof of $\boxed{l_2(V) \subset \text{Ran } Y_0^{1/2}}.$

Let $x \in l_2(V)$ and set

$$\begin{aligned} c &:= \|Vx\|^2 + \|V^2x\|^2 + \|V^3x\|^2 + \dots \\ P &:= \text{orthoprojection mapping onto } \text{span}\{x\} \end{aligned}$$

Then

$$\langle x, Y_t^{-1}x \rangle = \langle x, \frac{1}{t}Q_tx \rangle \leq \frac{1-z}{t}c = \frac{2c}{1+t} \leq 2c$$

Hence

$$\|x\|^2 P \leq 2cY_t, \quad \|x\|^2 P \leq 2cY_0, \quad \text{Ran } P \subset \text{Ran } Y_0^{1/2}$$

□

Remark to this proof.

- (i) $Y_t \gg 0, Y_0 \geq 0, P = P^{1/2} \geq 0$
- (ii) $\|Ax\| \leq \|Bx\| \quad (x \in H) \Rightarrow \text{Ran } A^* \subset \text{Ran } B^*$
(A and B are bounded). □

Corollary. $\boxed{l_2(V) = \text{Ran } Y_0^{1/2}}.$

3 Nonspectral Decomposition

Observation 1.

It follows from the definitions of X_t, Y_t that $X_t Y_t = Y_t X_t = t^2$. Hence $X_0 Y_0 = Y_0 X_0 = 0$ and $\overline{\text{Ran } X_0} \subset \text{Ker } Y_0$, $\overline{\text{Ran } Y_0} \subset \text{Ker } X_0$. But X_0, Y_0 are selfadjoint. Thus we obtain an orthogonal decomposition

$$H = \overline{\text{Ran } Y_0} + (\text{Ker } X_0 \cap \text{Ker } Y_0) + \overline{\text{Ran } X_0}$$

such that

- 1) first component is V -surinvariant;
- 2) third component is V^{*-1} -surinvariant.

(recall, some L is said to be T -surinvariant, iff $TL = L$). Moreover, denote $j_t := (I + tQ_t^{-1})^{1/2} / (I + tQ_t)^{1/2}$, then j_t is uniformly positive, bounded, there hold

$$(1 + \|V\|^2)^{-1/2} \leq j_t \leq (1 + \|V^{-1}\|^2)^{1/2},$$

and $(j_t V)^* Q_t (j_t V) = Q_t$. In particular $j_t V$ is similar to a unitary operator.

For a moment suppose $\dim H < \infty$. It is clear that now the restriction of V onto $\overline{\text{Ran } Y_0}$ and the restriction of V^{*-1} onto $\overline{\text{Ran } X_0}$ are similar to uniform contractions (see theorems 2.1 with remarks to the proof of $l_2(V) = \text{Ran } Y_0^{1/2}$). In addition, if

$$\text{Ran } Y_0 = \{0\} = \text{Ran } X_0,$$

then the restriction of V onto $\text{Ker } X_0 \cap \text{Ker } Y_0$ has the unit spectrum. $\square$

This motivates the

Definition 1. We shall say, a linear bounded operator T is **near similar to uniform contraction**, iff there exists a bounded operator $Y > 0$ such that

$$T Y T^* = \frac{Y}{I + Y}$$

We shall say, T is **s-approximately similar to a unitary**, iff there exists a net $\{j_t\}_t$ of bounded uniformly positive operators such that

- 1) $1/M \leq j_t \leq M$ for some real M (and every t),
- 2) $\text{strong} - \lim j_t = I$ (hence $\text{strong} - \lim j_t^{-1} = I$),
- 3) for every fixed t the operator $j_t T$ is similar to a unitary operator. $\square$

With this definition one can resume the section as follows:

Theorem 1. There exists an orthogonal decomposition

$$H = H_{<} + H_{=} + H_{>}$$

such that

- 1) $H_{<}$ is V -surinvariant and the restriction of V onto $H_{<}$ is near similar to a uniform contraction;
- 2) $H_{>}$ is V^{*-1} -surinvariant and the restriction of V^{*-1} onto $H_{>}$ is near similar to a uniform contraction;
- 3) If $H_{<} = \{0\} = H_{>}$ then V is s-approximately similar to a unitary;
- 4) $H_{<} \subset \overline{l_2(V)}$, $H_{>} \subset \overline{l_2(V^{*-1})}$.

Proof. Set $H_{<} := \overline{\text{Ran } Y_0}$, $H_{=} := \text{Ker } X_0 \cap \text{Ker } Y_0$, $H_{>} := \overline{\text{Ran } X_0}$ and apply the text. $\square$

4 Stability of U in terms of Q_t

Proposition 1. *Let*

$$\frac{1}{M}I \leq Q_t \leq MI$$

for some number $M > 0$ (and every t). Then V is similar to an unitary operator.

Proof. Let Q_0 be a weak limit point of the net $Q_t, t \rightarrow +0$. It exists and

$$\frac{1}{M}I \leq Q_0 \leq MI$$

$$Q_0 = V^*Q_0V$$

(see Observation 2.2). Hence, V is similar to an unitary operator. □

Proposition 2. *Let V be similar to an unitary operator. Then*

$$\frac{1}{M}I \leq Q_t \leq MI$$

for some number $M > 0$ (and every t).

Proof. For assumed V there exists a number $M > 0$ such that for every natural n there hold

$$\|V^n\|^2 \leq M, \quad \|(V^*)^{-n}\| \leq M$$

Take arbitrary $x \in H$, number $z, 0 < z < 1$ and apply the Observation 2.3:

$$\begin{aligned} \langle x, Q_t x \rangle &\leq (1-z)(\|Vx\|^2 + z\|V^2x\|^2 + \dots + z^{n-1}\|V^n x\|^2 + \dots) \\ &\leq (1-z)\frac{M}{1-z} \langle x, x \rangle = M \langle x, x \rangle \end{aligned}$$

Hence, $Q_t \leq MI$.

Now apply the Observation 2.3 to Q_t^{-1} and V^{*-1} :

$$\begin{aligned} \langle x, Q_t^{-1} x \rangle &\leq (1-z)(\|V^{*-1}x\|^2 + z\|(V^{*-1})^2x\|^2 + \dots) \\ &\leq M \langle x, x \rangle \end{aligned}$$

Hence, $Q_t^{-1} \leq M$ and $\frac{1}{M} \leq Q_t$. □

5 When Spectrum has Dichotomy.

Return us to the Example 1.1, which was called motivating . We remarked there, that R_0 is reminiscent of one of the spectral projector of the operator V (this operator was taken there to be normal).

Now we will show that somewhat similar situation holds always, especially when the spectrum of the operator V does not intersect the unit circle.

Observation 1. $\boxed{Y_0 = Y_0 R_0 = R_0 Y_0 = R_0^{1/2} Y_0 R_0^{1/2}}$ In addition R_0 acts on $Ran Y_0$, hence on $\overline{l_2(V)}$, as identity operator

Proof. Recall $Y_t = tQ_t^{-1}$, $Y_0 = s - \lim_{t \rightarrow +0} Y_t$, $R_t = (I + Q_t)^{-1}$, $R_0 \in w - \lim pt(R_t)$, all these operators are selfadjoint. What is more, the straightforward calculation shows that

$$Y_t(1 - R_t) = \frac{t}{I + Q_t} = (1 - R_t)Y_t.$$

The rest is obvious. □

Theorem 1. Suppose there is an V -invariant subspace, L say, such that $|\text{spectrum}(V|L)| < 1$; let P denote orthoprojector onto L ,
Then

a)

$$L \subset Ran Y_0^{1/2} \subset Ker(I - R_0)$$

b)

$$\|(I - R_t)P\| \rightarrow 0 \quad (t \rightarrow +0)$$

Proof.

a) Note $L \subset l_2(V)$ and apply theorem 2.1.

b)

Return to the Observation 2.3:

$$0 \leq Q_t \leq (1 - z)[V^*V + zV^{*2}V^2 + \dots + z^{n-1}V^{*n}V^n] + z^n V^{*n} Q_t V^n$$

Let V_P denote $PV|L$. Since $VP = PVP$, we can deduce that

$$0 \leq PQ_t P \leq (1 - z)[V_P^* V_P + zV_P^{*2} V_P^2 + \dots + z^{n-1} V_P^{*n} V_P^n] + z^n V_P^{*n} PQ_t P V_P^n$$

Next we adopt the spectrum argument. We make it in the same manner that standard practice suggests: there are some real positive ϵ , M such that

$$r(V_P) + \epsilon < 1, \quad \|V_P^n\| \leq M(r(V) + \epsilon)^n \quad (n = 1, 2, \dots),$$

Hence

$$\|z^n V_P^{*n} PQ_t P V_P^n\| \rightarrow 0 \quad (n \rightarrow \infty).$$

and it is routine matter to verify that

$$\|PQ_t P\| \leq \frac{(1 - z)}{1 - z(r(V_P) + \epsilon)^2} M^2(r(V_P) + \epsilon)^2.$$

Note that M and ϵ does not depend on z . So, we obtain $\|PQ_t P\| \rightarrow 0$ for $t \rightarrow +0$.

(One can show moreover : the serie

$$(1 - z)[V_P^* V_P + z V_P^{*2} V_P^2 + \cdots + z^{n-1} V_P^{*n} V_P^n \cdots]$$

is norm-convergent.)

Now note that

$$P R_t P \equiv P(I + Q_t)^{-1} P \geq P(I + P Q_t P)^{-1} P \xrightarrow{\parallel} 0$$

$$P(I - R_t)P \equiv P Q_t (I + Q_t)^{-1} P \leq P Q_t P (I + P Q_t P)^{-1} P \xrightarrow{\parallel} 0$$

Recall

$$0 \leq I - R_t \leq I$$

Hence

$$0 \leq P(I - R_0)^2 P = P(I - R_0)^{1/2} (I - R_0) (I - R_0)^{1/2} P \leq P(I - R_0) P$$

So, we can now establish that

$$\|(I - R_t)P\|^2 = \|P(I - R_t)(I - R_t)P\| \leq \|P(I - R_t)P\| \rightarrow 0,$$

It was to be proved. □

Corollary. *Suppose that the spectrum of the operator V does not intersect the unit circle; let P denote orthoprojector onto spectral subspace L corresponded to the set $\text{spectrum}(V) \cap \{z \in \mathbf{C} \mid |z| < 1\}$.*

Then

$$R_0 = P.$$

Proof. By Theorem 1

$$(I - R_0)P = 0.$$

Hence

$$P = R_0 P.$$

Next note that the equation

$$Q = V^* \frac{Q + t}{I + tQ} V, \quad Q \geq 0, \quad 0 < t \leq 1 \quad (*)$$

is equivalent to the equation

$$Q^{-1} = V^{-1} \frac{Q^{-1} + t}{I + tQ^{-1}} V^{*-1}, \quad Q \geq 0, \quad 0 < t \leq 1. \quad (**)$$

(for details see [Ch1,2])

For a moment introduce for the (unique) solution of (*) a longer denotation:

$$Q_t(V).$$

It is straightforward to deduce now that

$$Q_t(V^{*-1}) = Q_t(V)^{-1}, \quad R_t(V) = I - R_t(V^{*-1}), \quad Y_t(V) = X_t(V^{*-1}) \dots \text{etc.}$$

Last recall the Standard Spectrum Theorems (see e.g. [RS]) and apply Theorem 1 to the operator V^{*-1} . Then obtain

$$R_0(I - P) = 0.$$

Hence

$$R_0 = R_0 P.$$

To complete the proof let compare the second displayed formula with the last one in the current period. □

References

- [Ch1] S. A. Chorošavin. Some Existence Theorems for Nontrivial Invariant Majorant. 17 pp, VINITI 21.03.89 N 1765-V89 (Russ. Ref. Jour. Math. 1989, ref-no 7B931)
- [Ch2] S. A. Chorošavin. A theorem of existence of neutral invariant subspace. 9 pp, VINITI 06.07.89 N 4495-V89 (Russ. Ref. Jour. Math. 1989, ref-no 11B799)
- [Ch3] S. A. Chorošavin. Convergence of Angle Operators in case of Krein Approximations // (Voronezh Spring Mathematical School, 1996) Pontrjaginskije čtenija-7, Voronezh, 17-23 apr., 1996, p 181 (Russ. Ref. Jour. Math. 1996, ref-no 11B824)
- [Ch4] S. A. Chorošavin Some Decomposition of Linear Bounded Invertible Operator. // (Voronezh Spring Mathematical School, 1997) Pontrjaginskije čtenija-8, Voronezh, 4-9 maja, 1997, p 159 (Russ. Ref. Jour. Math. 1997, ref-no 10B706)
- [Krein64] M. G. Krein : A New Application of the Fixed-Point Principle in the Theory of Operators in a Space with Indefinite Metric. //DAN SSSR. 1964. **154**, N 5, 1023 –1026.(russisch)
- [Dal-Kre] Ju. L. Daletskij, M. G. Krein *The Stability of the Solutions of Differential Equations in a Banach Space*, Moscow.: Nauka, 1970 [Russian]
- [Krein65] M. G. Krein *Introduction to the geometry of indefinite J -spaces and to the theory of operators in those spaces*. In: Second mathematical summer school, Part 1, pp 15-92, Kiev.: Naukova dumka, 1965 [Russian]
- [Ch97] Chorošavin S. A. On one M. G. Krein problem. //TRANSACTIONS of RANS, series MMMIC, 1997, v.1, N.2, 95-101.
- [Ch98] Chorošavin S. A. An Example of J -Unitary U wich Has no Nonzero Invariant Subspace L such that $r(U|L) \leq 1$. //TRANSACTIONS of RANS, series MMMIC. 1998. **v.2**, N 2, 97–103
- [RS] M. Reed , B. Simon, *Functional analysis*, Methods of Modern Mathematical Physics, vol 1, - N.Y.: Academic Press, 1972.